
Clupeonella

()

caspia (Svetovidov, 1941)

*

(/ / : / / :)

Clupeonella caspia

($L_{\infty}=128.7\text{mm}$, $k=0.41\text{yr}^{-1}$, $t_0=-0.59\text{yr}$) $(R^2=0.918) \quad W=0.000001FL^{2.92}$

/ ± /

GSI

/ yr⁻¹

/

/ : : : /

/ yr⁻¹ / yr⁻¹

Clupeonella caspia :

(Fazli *et al.*, 2007; Parafkandeh

Haghghi, 2009)

.(Karimzadeh, 2011)

(Pourgholam *et
al.*, 1996)

(*C. Clupeonella caspia*, Svetovidov, 1941)

(*C. engrauliformis*, Borodin, 1904)

(Kazanchev, *grimmi*, Kessler, 1877)

M.

.1981; Berg, 1948)

leidyi

.(Melinikov, 2000)

(Fazli, 1990; Aseinova,
1992; Razavi Sayyad, 1993; Pourgholam *et al.*,
1996)

.(Mamedov, 2006; Shariati, 1994)

(Fazli *et al.*, 2007;

Mnemiopsis .Karimzadeh *et al.*, 2010)

(Ivanov *et al.*, *leidyi*
2000)

(Fazli *et al.*, 2004;

.Aliasghari, 2009)

)

.

(Berg, 1948)

(Besharat

and Khatib, 1993; Razavi Sayyad, 1993)

(Pourgholam *et al.*, 1996; Fazli *et al.*,

(Karimzadeh *et al.*, 2010 2002)

Fatemi *et al.*, 2009; Parafkandeh Haghghi, 2009;
Fazli *et al.*, 2002, 2004, 2005, 2007; Sayyad

Bourani, 1997)

mm

g

()

(Zhang and E = $\frac{F}{F + M}$.
Sullivan, 1988)

Microsoft Office Excel 2003 SPSS 10.0
chi-square FISAT

(Campana
.and Neilson, 1985; Francis and Campana, 2004)

$$W = a L^b$$

a (mm) L (g) W
(Bagenal,
t b
t=(b-3)/Sb .1978)
Sb b (Morey *et al.*, 2003)

% /
% /
% /
()
/ ± / mm / ± / g

$$L_t = L_\infty [1 - \exp^{-k(t-t_0)}]$$

$$R^2=0.918$$

$$W = 0.000001 FL^{2.92}$$

/ b ()
(b<3)
t
(p<0.05)

$$GSI = \frac{w}{W} \times 100$$

GSI .(Biswas, 1993)
(g) W (g) w
(S) (Z)
(Ricker, 1975)
Z = -\ln S
(M) (Z)

(L_∞=128.7mm,

$$k=0.41 \text{yr}^{-1}, t_0=-0.59 \text{yr}$$

$$() L_t = 128.7 [1 - \exp^{-0.41(t+0.59)}]$$


$$\log(M) = -0.0066 - 0.279 \log(L_\infty) + 0.6543 \log(K) + 0.4634 \log(T)$$

T


(Prakarn, 2002)

F=Z-M (F)

/	/	/	/	/	/
/	/	/	/	/	/
/	/	/	/	/	/




()



()

· · · () () / ± / / ± / .
 () (p<0.05) () () / ± / ()



()

p
/ / / / / / / / / / / / / / / / / / / /


$$\text{GSI} \quad .(\quad) \\ .(\quad) \quad / \pm / \\ / \quad \text{yr}^{-1}$$

/ yr⁻¹

$$| \quad \text{yr}^{-1} \quad | \quad \text{yr}^{-1}$$

/

(N=) ()



b

/

(Fazli *et al.*, 2007) /

(Karimzadeh *et al.*, 2010) /

(Salmanov, 1999; Ivanov, 2000)

b

(b<3)

(p<0.05)

t

(Vinogradov *et al.*, 1998;
Volovik, 2000; Kuliyev, 2004)

(Karimzadeh, 2011)

$t_0 = -0.59 \text{ yr}$ $k = 0.41 \text{ yr}^{-1}$ $L_\infty = 128.7 \text{ mm}$

L_∞

/

(Fazli, 2007)

(k)

/ yr⁻¹

(Janbaz

(Fazli, 2007)

(Karimzadeh *et al.*, 2010)

and Abdolmaleki, 2008)

/ yr⁻¹ / yr⁻¹ *et al.*, 2010

/

(Fazli *et al.*, 2007)

(Karimzadeh *et al.*, 2010)

/

(Fazli *et al.*, 2007)

(Karimzadeh *et al.*, 2010)

/

(Stevensen and Campana,
1992)

L_∞

() ()

(Ben-Yami, 1976)

(Maeda,

.1951)

(Fazli *et al.*,

.2007)

(Karimzadeh *et al.*, 2010)

/

(Ben-Yami, 1976)

)

(

(Wootton, 1990)

-

(Prikhodko,

.1981)

/ :

(Razavi Sayyad, 1993)

(Pourgholam *et al.*, 1996)

(Prikhodko, 1981)

(Abtahi *et al.*, 2002)

(Fazli *et al.*,

(Fazli *et al.*, 2005)

.2007)

(Karimzadeh *et al.*, 2010)

(Karimzadeh

: /

et al., 2010)

(Beverton and Holt, 1959)

/ yr⁻¹

/ yr⁻¹

(Kideys *et al*, 2005;

(/ yr⁻¹)

yr⁻¹

M. leidyi .Kideys and Moghim, 2003)

.(Kideys *et al.*, 2001)

(Fazli *et al.*, 2007)

(Rouhi, 2003)

(Karimzadeh, 2011)

/ yr⁻¹

(Roshantabari *et al.*,

.2003; Mirzajani *et al.*, 2005)

/ yr⁻¹ / yr⁻¹

()

(Fazli *et al.*, 2005)

(Janbaz *et al.*, 2010)

(Fazli *et al.*,

.2007; Karimzadeh *et al.*, 2010)

(Parafkandeh, 2009)

1

(Gulland, 1983)

(E \approx .)

(Fazli *et al.*, 2007)

(Karimzadeh *et al.*, 2010)

References

- Abtahi, B., Taghavi, H., Fazli, H., Usefian, M., 2002. Study of fecundity and biometrical characteristics of common kilka (*Clupeonella delicatula*) in Mazandaran province (Babolesar). Iranian Journal of Marine Sciences 1(2), pp1-9 (In Persian).
 - Aliasghari, M., 2009. Aquatic invertebrates. Avaye Masih Pub. ISBN: 978-964-2769-42-1. 338p (In Persian).
 - Aseinova, A.A., 1992. Common Kilka in the Caspian Sea. Ichthyo fauna and commercial resources. Nauka. Moscow. pp 71-80.
 - Bagenal, T.B., 1978. Methods of assessment of fish production in freshwater. Blackwell scientific publication. 365p.

...

- Bagheri, S., Sabkara, D., Roohi, A., Parafkandeh, F., Ghasemi, Sh., Razavi Sayyad, P., 2005. Studying on distribution and abundance of Ctenophora in southern Caspian Sea. Final report. Iran Fisheries Research Institute. 32 p (In Persian).
- Bagheri, S., Mirzajani, A., Sabkara, D., Makaremi, M., Roohi, A., Khodaparast, H., Malekshomali, M., Ghasemi, Sh., Parafkandeh, f., 2007. Studying on feeding and reproduction of Ctenophora in southern Caspian Sea. Final report. Iran Fisheriees Research Institute. 37 p (In Persian).
- Ben-Yami, M., 1976. Fishing with light. FAO of the united nations, Fishing News Books. 121p.
- Berg, L.S., 1948. Freshwater fishes of the U.S.S.R. and adjacent countries. Vol: 1, 2, 3, pp1-223.
- Besharat, K., Khatib, S., 1993. Determination of kilka (genus: *Clupeonella*) fishing area in northern parts of Iran and hydrologic and hydrobiologic stydies on the Caspian Sea. Final Report. Mazandaran Fisheries Research Center. 181 p (In Persian).
- Beverton, R.J.H., Holt, S.J. 1959. A review of the life spans and natural mortality rates of fish in nature and their relation to growth and other physiological characteristics. Central Institute of Brackishwater Aquaculture colloquium on ageing Vol 5, pp 142-180.
- Biswas, S.P., 1993. Manual of Methods in fish biology. South Asian Publishers, PVT Ltd, New Delhi, India. 157p.
- Campana, S.E., Neilson, J.D., 1985. Microstructure of fish otoliths. Canadian Journal Fish Aquatic Science 42, pp 1014-1032.
- Fatemi, S.M.R., Kaymaram, F., Parafkandeh Haghghi, F., Vosooghee, G.H., Taghavi Motlagh, S.A., 2009. Validation of back-calculation methods using otoliths to determine the length of anchovy kilka (*Clupeonella engrauliformis*). Iranian Journal of Fisheries Sciences 8(2), pp115-126 (In Persian).
- Fazli, H., 1990. Biology of genus *Clupeonella* in Caspian Sea. Conference of suitable exploitation of the Caspian Sea aquatics. September 1990. Babolsar. Pp 81-89.
- Fazli, H., 2007. Population Dynamics and Stock Assessment of Kilka (Genus: *Clupeonella*) in Iranian Waters of the Caspian Sea. Phd thesis, Pukyong National University, Department of Marine Biology. 125p. Available on: <http://kosfc.chonnam.ac.kr>
- Fazli, H., Rouhi, A., 2002. Probable effect of ctenophore entrance on kilka species composition, catch and stock in southern Caspian Sea (1997-2001). Iran Fisheries Journal No1, 63-72 (In Persian).
- Fazli, H., Bourani, M., Janbaz, A., Naderi, M., Abo, M., Moghim, M., Ofi, F., Azari, A., 2002. Study of kilka biology and statistics in commercial fishing regions. Mazandaran Fisheries Research Center. 112pp (In Persian).
- Fazli, H., Sayyad Bourani, M., Janbaz, A.A., 2004. Bigeye kilka (*Clupeonella grimmi*) biological characteristics in Iranian commercial fishing during 1997-2001. Iran Fisheries Journal 13(4), pp125-138 (In Persian).
- Fazli, H., Sayyad Bourani, M., Janbaz, A.A., 2005. Common kilka (*Clupeonella cultriventris caspia*) biological characteristics and the effects of *Mnemiopsis Leidyi* in southern part of the Caspian Sea. Researching and Making Journal (aquatics) 69, pp87-96.
- Fazli, H., Zhang, C. I., Hay, D. E., Lee, C. W., Janbaz, A. A., Borani, M.S., 2007. Population Dynamics and Stock Assessment of Common Kilka (*Clupeonella cultriventris caspia*) in the Caspian Sea. Iranian Journal of Fisheries Sciences 7(1), pp47-70.
- Fazli, H., Zhang, C.I., Hay, D.E. and Lee, C.W., 2009. Multi-species approach for stock management of kilka fish (genus: *Clupeonella*) in Iranian waters of the Caspian Sea. Iranian Journal of Fisheries Sciences 8 (2), pp141-154.
- Francis, R.C., Campana, S.E., 2004. Inferring age from otolith measurements: a review and a new approach. Canadian Journal of Fish 61, pp1269-1284.
- Gulland, J.A. 1983. Fish stock assessment. A manual of basic methods. Chichester John Wiley. FAO. Wiley Series on Food and Agriculture Vol. 1, 223p.
- Ivanov, P.I., 2000. Biological resources of the Caspian Sea. KaspNirkh, Astrakhan. 130p.
- Ivanov, P.I., Kamakim, A.M., Ushivtzev, V.B., Shiganova, T.A., Zhukova, O., Aladin, N., Wilson, S.I., Harbinson, G.R., Dumont, H.J., 2000. Invasion of the Caspian Sea by the comb jellyfish *Mnemiopsis leidyi* (Ctenophora). Journal of biological invasion 2, pp255-258.
- Janbaz, A., Abdolmaleki, Sh., 2008. Age , growth and mortality of clupepnella cultriventris caspia in the coastal mazandaran. Pajouhesh And Sazandegi, Animal and Fisheries Sciences, 21(81), pp 78-80 (In Persian).

-
- Janbaz, A., Khedmati, K., Shabani, Kh., Fazli, H., Parafkandeh, F., Abdolmaleki, S., Moghim, M., Afraei, M., Daryanabard, Gh., 2010. Project of studying the kilka fishes biological parameters (age, growth, feeding, reproduction) in southern Caspian Sea. Iranian Fisheries Research Organization. 112 p (In Persian).
 - Jones, R., 1981. Use of length composition data in fish stock assessment. FAO fisheries circulation. 734, FAO. Rome, Italy, 55 p.
 - Karimzadeh G., Gabrielyan, B., Fazli, H., 2010. Population dynamics and biological characteristics of kilka species (Pisces: Clupeidae) in the southeastern coast of the Caspian Sea. Iranian Journal of Fisheries Sciences 9(3), pp422-433.
 - Karimzadeh, G., 2011. Study of natural and fishing mortality and exploitation rate of common kilka *Clupeonella cultriventris* in southeast part of the Caspian Sea (Babolsar). Annual Review and Research in Biology 1(2), pp33-43.
 - Kazanchev, A., 1981. Caspian Sea and its basin fishes. Translator: Shariati, A., (1991). Iran Fisheries Publication. 171p.
 - Kideys, A.E., Shahram, G., Davood, G., Roohi, A., Bagheri, S., 2001. Strategy for combating *Mnemiopsis* in the Caspian waters of Iran. Final report, July 2001, prepared for Caspian environment program, Baku, Azerbaijan. 2001.
 - Kideys, A.E., Moghim, M., 2003. Distribution of the alien ctenophore *Mnemiopsis leidyi* in the Caspian Sea in the August 2001. Marine Biology 142, pp 163-171.
 - Kideys, A.E., Roohi, A., Bagheri, S., Finenko, G., Kamburska, L., 2005. Impacts of invasive ctenophores on the fisheries of the Black Sea and Caspian Sea. Oceanography 18(2), pp76-85.
 - Kuliyev, Z., 2004. Dynamics of *Mnemiopsis leidyi* distribution in the Azerbaijan sector of the Caspian Sea in 2001- 2002. In Aquatic Invasion in the Black, Caspian and Mediterranean Seas, pp 203-204.
 - Maeda, H., 1951. Analytical studies on marine lamp-communities. Publication of the Seto Marine Biological Laboratory 1(4), pp195-213.
 - Mamedov, E.V., 2006. The biology and abundance of kilka (*Clupeonella* spp.) along the coast of Azerbaijan, Caspian Sea. ICES Journal of Marine Science 63, pp 1665-1673.
 - Melnikov, V.N., 2000. Kilka fishing methods with funnel-suction net. Caspian Sea Bony Fish Research Center. 24 P.
 - Mirzajani, A., Bagheri, S., Sabkara, J., Khodaparast, H., Malekshomali, M., Parafkandeh, F., Ghasemi, Sh., Negarestan, H., 2005. Studying on controlling the Caspian Sea Ctenophora. Final report. Iran Fisheries Research Institute. 69 p.
 - Morey, G., Moranta, J., Massuti, E., Grau, A., Linde, M., Riera, F., Morales-Nin, B., 2003. Weight-length relationships of littoral to lower slope fishes from the Western Mediterranean. Fisheries Research 62, pp 89-96.
 - Parafkandeh, F., 2009. Population dynamics of kilka fishes in southern Caspian Sea. PhD thesis in marine biology. Islamic Azad University, Sciences and Researches Branch. 117p.
 - Pauly, D., 1984. Length-converted catch curves. A powerful tool for fisheries research in the tropics (Part III). International Center for Living Aquatic Resources Management Fishbyte 2(3), pp9-10.
 - Pauly, D., 1999. On interrelationships between natural mortality, growth parameters and mean environment temperature in 175 fish stock. J. Cons. Calcutta Institute of Engineering and Management 39, pp175-192.
 - Prakarn, S., 2002. Building Awareness in Aspects of Fishery Statistics, Stock Assessment and Management. Proceedings of the FAO/SEAFDEC Regional Training Workshop on the Use of Statistics and Other Information for Stock Assessment. RAP PUBLICATION. 108 p.
 - Prikhodko, B.I., 1981. Ecological features of the Caspian kilka (genus *Clupeonella*). Scripta Publishing Co. pp 27-35.
 - Pourgholam, R., Sedov, V., Yermalchev, V., Besharat, K., Fazli, H., 1996. Stock assessment of kilka fishes by hydro acoustic method, 1994-1995. Final report, Mazandaran Fisheries Research Center. 125 p (In Persian).
 - Razavi Sayad, B., 1993. Abundance and distribution of kilka in Iranian waters. Scientific bulletin of Iran Fisheries No 2, pp 11-25.
 - Ricker, W.E., 1975. Computation and interpretation of biological statistics of fish populations. Bulletin of Fisheries Research Board of Canada 191, pp1-382.
 - Roshantabari, M., Takmilian, K., Sabkara, J., Rouhi, A., Rostamian, M., 2003. Zooplankton distribution in southern Caspian Sea. Iranian Scientific Fisheries Journal 2(3), pp 83-96 (In Persian).

...

-
- Rouhi, A., 2003. Study and distribution of ctenophores and the possibility of biological battle in southern Caspian Sea. Mazandaran Fisheries Research Center. 60 p (In Persian).
 - Rouhi, A., Yasin, Z., Kideys, A., Hwai, A.T.S., Ganjian Khanari, A., Develi, E., 2008. Impact of a new invasive ctenophore (*Mnemiopsis leidyi*) on the zooplankton community of the Southern Caspian Sea. *Marine Ecology* 29(4), pp 421-434.
 - Salmanov, M.A., 1999. Ecology and biological reproduction of the Caspian Sea. Edited by: U.I. Sorokin. Baku. 1999. 397p.
 - Sayyad Bourani, M., 1997. Survey on some biological characteristics of anchovy kilka (*Clupeonella engrauliformis*) in Guilan waters. *Iran Fisheries Journal* 6(1), pp59-70.
 - Shariati, A., 1994. Animals and biological production in Caspian Sea. Fisheries research center of Guilan Province. 405 p.
 - Stevenson, D.K., Campana, S.E., 1992. Otolith microstructure examination and analysis. Canadian Special Publication of Fisheries and Aquatic Science Issue117, 126p.
 - Vinogradov, M.E., Shushkina, E.A., Musaeva, E.I., Sorokin, P.Y., 1998. Ctenophore *Mnemiopsis leidyi* (A. Agassiz Ctenophora: Lobata). *Newsletter in the Black Sea. Oceanology* (29), pp 293-298.
 - Volovik, S.P., 2000. The Ctenophore *Mnemiopsis leidyi* (A. Agassiz) in the Azov and Black Seas: Biology and Consequences of its Intrusion. Rostov-on-Don, 2000. 498p
 - Wootten, R.J., 1990. Ecology of teleost fishes. Chapman and Hall, London, UK. 404p.
 - Yermalchev, V., Sedov, S.A., 1990. Caspian Sea kilka fishes stock assessment with hydroacoustic method. Murmansk. Russia. 90 p.
 - Zhang, C.I., Sullivan, P.J., 1988. Biomass-based cohort analysis that incorporates growth. *Transactions of American Fisheries Society* 117, pp 180–189.

Investigation of Some of the Growth and Reproductive Parameters of Common Kilka *Clupeonella caspia* (Svetovidov, 1941) in Southern Caspian Sea (Mazandaran Zone)

M. Aliasghari^{*1}, A. Keramat Amirkolaie² and F. Parafkandeh Haghghi³

¹ Young Researchers Club, Qaemshahr Branch, Islamic Azad University, Qaemshahr, I.R. Iran

² Departments of Fisheries, Sari Agricultural and Natural Resources University, Sari, I.R. Iran

³ Iran Fisheries Research Organization, Tehran, I.R. Iran

(Received: 30/07/2011 , Accepted: 09/01/2012)

Abstract

The present research aimed to study the growth and reproductive parameters of common Kilka *Clupeonella caspia* in southern Caspian Sea in 2010. After measuring the length and weight of 3774 samples, they were dissected, sex determined and female gonads were weighed. Sexual maturity classification was carried out based on six stages. Age was determined using sagitta otoliths. Length-weight relationship was $w = 0.000001FL^{2.92}$ ($R^2=0.918$). Growth parameters were ($L_\infty=128.7\text{mm}$, $k=0.41\text{yr}^{-1}$, $t_0=-0.59\text{yr}$). Age composition was constituted of 6 age groups, including 1-6 years. Age average was 3.27 ± 0.09 years. 3 years old fish were the largest age group, composing 45.24% of the catch population. Male: female ratio was 1: 0.779 and males were dominant. Study of sexual maturity stages and GSI showed that common Kilka spawning begins in February and the peak was in May and June. Survival rate was 0.239 yr^{-1} . Natural and fishing mortality coefficients were 0.448 yr^{-1} and 0.983 yr^{-1} . The exploitation rate calculated up to 0.687 indicating the species over fishing. Our results showed that the mean length and weight of common Kilka population has increased but the mean age decreased during the last years.

Keywords: *Clupeonella caspia* ; Caspian Sea; Growth; Reproduction; Age

*Corresponding author: Tel: +981112145223 Fax: +981112145223 E-mail: aliasghari_mehrdad@yahoo.com